

Extensive Disease Small Cell Lung Cancer Dose-Response Relationships

Implications for Resistance Mechanisms

David J. Stewart, MD, FRCPC,* Constance Johnson, MD, PhD,¶ Adriana Lopez, MS,†
Bonnie Glisson, MD,* Jay M. Rhee, MD,‡ and B. Nebiyu Bekele, PhD†

Background: Some studies (but not others) suggested that high doses are beneficial in small cell lung cancer (SCLC). We hypothesized that dose-response curve (DRC) shape reflects resistance mechanisms.

Methods: We reviewed published SCLC clinical trials and converted response rates into estimated mean tumor cell kill, assuming killing is proportional to reduction in tumor volume. Mean % cell survival was plotted versus planned dose intensity. Nonlinear and linear meta-regression analyses (weighted according to the number of patients in each study) were used to assess DRC characteristics.

Results: Although associations between dose and cell survival were not statistically significant, DRCs sloped downward for five of seven agents across all doses and for all seven when lowest doses were excluded. Maximum mean cell kill across all drugs and doses was approximately 90%, suggesting that there may be a maximum achievable tumor cell kill irrespective of number of agents or drug doses.

Conclusions: Downward DRC slopes suggest that maintaining relatively high doses may possibly maximize palliation, although the associations between dose and slope did not achieve statistical significance, and slopes for most drugs tended to be shallow. DRC flattening at higher doses would preclude cure and would suggest that “saturable passive resistance” (deficiency of factors required for cell killing) limits maximum achievable cell kill. An example of factors that could flatten the DRC at higher doses and lead to saturable passive resistance would be presence of quiescent, non-cycling cells.

Key Words: Small cell, Dose-response, Resistance, Quiescence.

(*J Thorac Oncol.* 2010;5: 1826–1834)

*Department of Thoracic/Head and Neck Medical Oncology, †Department of Biostatistics, and ‡Division of Cancer Medicine, University of Texas M. D. Anderson Cancer Center, Houston, Texas; and ¶Duke University, Durham, North Carolina.

Disclosure: The authors declare no conflicts of interest.

Address for correspondence: David J. Stewart, MD, FRCPC, Department of Thoracic/Head & Neck Medical Oncology, University of Texas M. D. Anderson Cancer Center, 1515 Holcombe Blvd, Unit 432, Houston, TX 77030. E-mail: dstewart@mdanderson.org

Copyright © 2010 by the International Association for the Study of Lung Cancer

ISSN: 1556-0864/10/0511-1826

Chemotherapy with or without radiotherapy for patients with limited disease (LD) small cell lung cancer (SCLC) has a 5-year survival rate of 10 to 25%.^{1,2} In extensive disease (ED) SCLC, chemotherapy gives complete responses in more than 20% of patients and prolongs median survival.^{1,2} Common regimens used in SCLC include etoposide-cisplatin (EP) and cyclophosphamide-doxorubicin-vincristine (CAV). Carboplatin, ifosfamide, epirubicin, paclitaxel, vinorelbine, topotecan, irinotecan, gemcitabine, pemetrexed, and other agents may also be used.^{1–5} Patients with prior responses often benefit from further chemotherapy if progression occurs more than 3 months after completing prior chemotherapy (“sensitive relapse”), whereas they will generally be resistant to further chemotherapy if progression occurs during or rapidly after completing front-line chemotherapy (“refractory relapse”).⁶ Although the major factors underlying resistance to chemotherapy remain undefined, numerous factors may contribute.^{7,8}

We postulated that dose-response curve (DRC) shape (log % cell survival versus dose) would reflect major resistance mechanisms limiting drug efficacy (Figure 1).^{8,9} By this hypothesis, excess of a resistance factor such as an efflux pump and DNA repair (“active resistance”) would give a DRC shoulder, analogous to competitive inhibition of drug effect. In this situation, the DRC slope would steepen after resistance factors had been overwhelmed by higher doses. One would not expect DRC flattening, because one would not expect the resistance mechanism to be ineffective at low doses and then become increasingly effective at higher doses. Mutation or alteration of a factor such as a target or drug uptake mechanism (“nonsaturable passive resistance”) would give reduced DRC slope, pharmacodynamically analogous to decreased affinity of a drug for its receptor. The rationale for the explanation of nonsaturable passive resistance is that one would not expect the efficacy of a resistance factor to increase proportionately to the effect of an increasing drug dose. Saturation or deficiency of something required for cell killing (e.g., a drug uptake system or cells in a sensitive phase of the cell cycle) (“saturable passive resistance”) would give DRC flattening at higher doses, analogous to noncompetitive inhibition of drug effect. In non-small cell lung cancer (NSCLC), we found that DRCs flatten at higher doses for all regimens assessed, suggesting that in NSCLC efficacy is limited by saturation or deficiency of factors required for drug effec-




FIGURE 1. We postulated that dose-response curve shape would reflect whether resistance was due to deficiency or saturation of a factor required for drug effect (which would result in flattening of the dose-response curve at higher doses, “saturable passive resistance,” analogous to noncompetitive inhibition of drug effect), due to mutation, etc., resulting in decreased affinity of a drug for an uptake system, target, etc. (which would give decreased curve slope, “nonsaturable passive resistance”), or due to excess of a resistance factor (which would give a shoulder on the dose-response curve if log effect is plotted against linear dose, “active resistance,” analogous to competitive inhibition of drug effect). Reprinted with permission from *Crit Rev Oncol Hematol*.⁸

acy.¹⁰ Herein, we used DRC assessments in SCLC as a basis for generating hypotheses regarding resistance mechanisms limiting drug efficacy in extensive disease.

METHODS

To be eligible for inclusion, a study had to report response rate using World Health Organization¹¹ or RECIST¹² criteria for front-line chemotherapy in ED SCLC. Studies were not included if they used alternating or cross-over designs, used radiotherapy, did not differentiate between ED and LD SCLC or between SCLC and NSCLC (if both LD and ED or both SCLC and NSCLC were included in the study), were restricted to tumor in a specific anatomic location (e.g., brain), or were testing primarily maintenance therapy or new drug administration methods or schedules. Using PUBMED and manual searches, we identified 46 studies (reporting 62 drug combinations) published between 1985 and 2005 that met all inclusion criteria (Table 1).^{13–58}

We used response rate to roughly estimate mean % tumor cell kill for each trial. We assumed tumor cell kill was proportional to reduction in tumor volume, recognizing that several factors would reduce estimate accuracy, including tumor regrowth between cycles, residual necrotic or stromal cells, tumor measurement inaccuracies, tumor shape irregularity, etc., and recognizing that the relationship between response rate and % tumor cell kill has to our knowledge never actually been proven in either preclinical or clinical investigations. For complete responses (CR), we assumed

more than 99% tumor cell kill. For a partial response, a tumor diameter decreases to less than 0.7 of the original diameter (with a decrease in the product of widest perpendicular diameters to $<0.7 \times 0.7 = 0.49$ of the original). This in turn translates roughly into a reduction in tumor volume to $<0.7 \times 0.7 \times 0.7 = 0.343$ of the original volume, such that the tumor has to have lost more than 65.7% of its original volume to be classified as a partial response. Hence, for the purposes of this study, we assumed that the average patient with a partial response would have an 83% reduction in tumor volume $[(100 + 65.7)/2]$ and an average 83% cell kill. Because stable disease (SD) represents a 30% reduction in tumor diameter up to a 12 to 20% increase in tumor diameter (depending on whether World Health Organization or RECIST criteria are used), we assumed that the average SD patient had a 13% cell kill. For patients with progressive disease (PD), we assumed a 0% cell kill, recognizing that this would substantially underestimate cell kill in some patients, such as those who had initial tumor shrinkage followed by rapid regrowth. Mean % tumor cell kill for the average patient in an individual study was calculated as $100 \times ([\text{no. CR} \times 0.99] + [\text{no. PR} \times 0.83] + [\text{no. SD} \times 0.13]) / \text{number of patients in the study}$.

Nonlinear meta-regression analysis (with each study weighted by number of patients) was then performed using SAS to model the relationship between mean % cell survival and dose intensity. Based on observations in our earlier studies in NSCLC,¹⁰ we hypothesized that in SCLC we would initially see a downward slope on the survival curve with

TABLE 1. Regimens Included

Regimen	No. of Studies
Cisplatin + etoposide	9
+ Ifosfamide	3
+ Paclitaxel	2
+ Irinotecan	2
+ Topotecan	1
+ Doxorubicin + vincristine	2
+ Doxorubicin + cyclophosphamide	1
+ Epirubicin + cyclophosphamide	2
+ Epirubicin + ifosfamide	1
Cisplatin + epirubicin	1
Cisplatin + irinotecan	2
Cisplatin + paclitaxel + topotecan	1
Carboplatin + etoposide	3
+ Vincristine	1
+ Epirubicin	2
+ Epirubicin + ifosfamide	1
Carboplatin + teniposide	1
Carboplatin + vinorelbine	2
Carboplatin + paclitaxel	1
+ Ifosfamide	1
Cyclophosphamide + epirubicin	1
Cyclophosphamide + etoposide	2
Etoposide	1
Etoposide + ifosfamide	1
Etoposide + paclitaxel	1
Etoposide + vincristine	1
Docetaxel	1
Docetaxel + gemcitabine	2
Doxorubicin + etoposide + ifosfamide	1
Epirubicin	1
Epirubicin + ifosfamide + vindesine	3
Paclitaxel	2
Topotecan	1
Vinorelbine	1

increasing dose, followed by a flattening of the curve at highest doses. To test this hypothesis, we assessed whether data could be fit using a weighted nonlinear regression function characterized by the following mean function:

% Tumor Cell Survival

$$= \gamma(\beta + (1 - \beta) \exp(\alpha \times \text{dose-intensity})),$$

subject to the restrictions $0 < \beta < 1$, $-5 < \alpha < 0$, $0 < \gamma < 100$, where the α parameter measures the relationship between dose and cell survival, β is the point at which cell survival plateaus, and γ is a scaling factor to permit scaling from 0 to 100%. Note that the α parameter characterizes the effect of dose on % cell survival and is set to be negative to ensure that % cell survival decreases with dose; the β parameter, as stated above, characterizes a level at which % cell survival plateaus and below which % cell survival does not fall. The plateau point characterized by β can be thought of as analogous to the cure rate fraction in some survival models.

TABLE 2. Number of Patients in Evaluable Studies

Drug	No. of Patients		
	Total No. of Patients (No. of Studies)	Study Size (Median)	Study Size (Range)
Cisplatin	2198 (30)	58	15–283
Cyclophosphamide	294 (8)	32	17–65
Carboplatin	605 (11)	43	28–156
Ifosfamide	588 (11)	54	16–87
Etoposide	2831 (40)	51	16–283
Epirubicin	680 (11)	65	30–78
Paclitaxel	575 (8)	42	34–283

Weighted linear regression analysis was used where data could not be fit to this nonlinear model. The criteria for using weighted linear regression models in place of weighted nonlinear models were when the weighted linear regression model resulted in lower (i.e., better) Akaike information criterion than the weighted nonlinear model. For carboplatin trials reporting dose as mg/m², AUC dose was estimated. For oral etoposide, 50% bioavailability was assumed.⁵⁹

Analyses were done over the entire dose range available for each drug and were repeated after eliminating lower doses at or below the lower limit of those commonly used.

RESULTS

Sufficient data were available to permit analyses for cisplatin, carboplatin, cyclophosphamide, ifosfamide, etoposide, epirubicin, and paclitaxel (Table 2), whereas data were insufficient for analysis of other agents. Figure 2 shows the curves over the entire dose range assessed for each drug. Data were fit using a weighted nonlinear meta-regression model for cisplatin and cyclophosphamide, whereas weighted linear regression models are presented for the other agents, because the β parameter could not be fit by the nonlinear regression model. The parameter estimates and SEs for the models when all data are used for modeling are given in Table 3. In addition, for each model, we present the fitted models in Figures 2 and 3. For each of cisplatin, cyclophosphamide, ifosfamide, etoposide, and paclitaxel, there was a downward slope on the DRC, with increasing cell kill with increasing dose intensity, although this association between dose intensity and cell kill did not achieve statistical significance. When lowest drug doses were omitted (Figure 3), data for each of cisplatin, carboplatin, cyclophosphamide, and ifosfamide could be fit by the weighted nonlinear meta-regression model, and there was a downward curve slope for all seven agents, although again, statistical significance was not achieved.

Similar results were obtained using nonweighted analyses and using log % cell survival rather than % cell survival (data not shown). Irrespective of drug doses and combinations used, mean cell kill generally did not exceed 90% (Figures 2 and 3). Although several studies achieved this cell kill, none substantially exceeded it no matter how many drugs were used and no matter what doses per cycle or total cumulative doses were used.




FIGURE 2. Weighted nonlinear meta-regression analysis (where the α parameter measures the relationship between dose and cell survival) or weighted linear regression analysis (where the β parameter measures the relationship between dose and cell survival) over the entire dose range assessed. Data for each of cisplatin ($n = 30$, $\alpha = -0.02$) and cyclophosphamide ($n = 8$, $\alpha = -0.01$) could be fitted to the nonlinear regression model, whereas data for each of etoposide ($n = 40$, $\beta = -0.01$), carboplatin ($n = 11$, $\beta = 0.14$), epirubicin ($n = 11$, $\beta = 0.25$), paclitaxel ($n = 8$, $\beta = -0.11$), and ifosfamide ($n = 11$, $\beta = -0.01$) could not be fitted to the nonlinear regression model and were instead assessed by linear regression analysis.

DISCUSSION

Despite often initially responding well to chemotherapy, ED SCLC remains incurable. The reasons for this incurability are unknown. Several resistance mechanisms have been described in SCLC cell lines and xenografts,^{8,60} but it is unclear which of these are important clinically. Although this study was limited by a relatively small number of evaluable studies and by our inability to correct for the contribution of concurrent agents in drug combination studies, the lack of a statistically significant association between dose intensity of any agent and estimated cell kill would argue against a steep DRC for any of the agents assessed. This is in keeping with our observation that there seems to be a maximum mean cell kill of around 90% across all ED SCLC studies, irrespective of drug type, number of drugs used, drug doses per cycle, dose intensity, total cumulative drug dose administered, treatment schedule, or cell cycle specificity of the agents used. This apparent maximum cell kill of around 90% irrespective of drug used would suggest

that DRCs for all agents flatten at higher drug doses, and that there is no combination, dose, or schedule of currently available agents that would be capable of curing ED SCLC. Even very high chemotherapy doses failed to improve outcome. This is similar to our observation of DRC flattening at higher doses in NSCLC¹⁰ and is in keeping with the inability of even very high drug doses with bone marrow transplantation to cure any metastatic epithelial malignancies.⁶¹ Overall, the observations in this study are in keeping with our earlier conclusion¹⁰ that there could possibly be a common factor that limits curability of all metastatic epithelial malignancies. If interpreted according to our hypothesis on the link between DRC shape and resistance mechanisms,⁹ this DRC flattening would suggest that the major problem is one of deficiency or saturation of a factor required for drug efficacy, rather than incurability being due to excess of resistance factors such as efflux pumps or DNA repair pathways. By our hypothesis,⁹ “active” resistance factors such as efflux pumps or DNA repair pathways would be most important at low drug doses

TABLE 3. Model Estimates, SEs, and 95% Confidence Intervals

Treatment	Model	Parameter	Parameter Estimate (SE)	Lower 95% Limit	Upper 95% Limit
Cisplatin	Nonlinear	α	-0.0233 (0.02)	-0.0624	0.0159
		β	0.00 (0.00)	0.000	0.000
		γ	63.36 (29.65)	2.63	124.10
Cyclophosphamide	Nonlinear	α	-0.0052 (0.003)	-0.011	0.00084
		β	0.0061 (0.200)	-0.484	0.4958
		γ	100.00 (0.00)	100.000	100.000
Carboplatin	Linear	Intercept	27.179 (11.730)	0.645	53.714
		Slope	0.137 (7.527)	-16.889	17.164
Ifosfamide	Linear	Intercept	43.636 (12.536)	15.277	71.994
		Slope	-0.006 (0.006)	-0.020	0.008
Etoposide	Linear	Intercept	35.671 (7.311)	20.871	50.471
		Slope	-0.014 (0.056)	-0.128	0.100
Epirubicin	Linear	Intercept	23.584 (10.728)	-0.685	46.754
		Slope	0.248 (0.334)	-0.507	1.003
Paclitaxel	Linear	Intercept	41.072 (30.349)	-33.189	115.332
		Slope	-0.105 (0.499)	-1.326	1.115

but should be surmountable by increasing drug doses. Although there was a downward slope on many of the DRCs, this was not statistically significant for any agent, and the observation of a maximum mean cell kill that was consistent across all agents suggests that there is an upper limit to the cell kill achievable by augmenting drug doses.

One possible explanation of the DRC flattening would be the presence of quiescent cells in the tumor that survive the first cycle of chemotherapy, irrespective of drug dose. Quiescent cells generally have increased resistance to therapy,^{62–64} and various preclinical observations support a link between quiescence and resistance. For example, resistance may be accompanied by decreased tumorigenic capacity,⁶⁵ a decrease in uptake across cell membranes of a broad range of factors and drugs,⁶⁶ downregulation of expression of various membrane receptors and transporters,⁶⁶ decreased amount and altered localization of a variety of membrane proteins,⁶⁵ faulty or decreased endocytosis,^{66,67} reversible senescence,^{68,69} autophagy,⁷⁰ or appearance of nondividing multinucleated cells that may later resume cell cycling.⁷¹

If quiescent cells did survive the first cycle of chemotherapy irrespective of dose (thereby flattening DRCs), these surviving quiescent cells might maintain their quiescent phenotype for a prolonged period of time after drug exposure but might ultimately start cycling again, as has been described for quiescent cells in some preclinical lung cancer systems.^{68,69,71} The fact that occasional late relapses are seen in patients with SCLC^{72–75} is in keeping with SCLC cells being capable of reversible quiescence. If resistance was due to quiescence, then the tumor cells might potentially be sensitive again when they ultimately resumed cycling. Such a pattern of prolonged quiescence followed by drug-sensitive tumor regrowth would be in keeping with the observation in SCLC that patients who do not require reinitiation of chemotherapy until more than 3 months after their prior therapy may again be sensitive to chemotherapy when they ultimately do suffer tumor regrowth (the “sensitive relapse” phenotype⁶). In addition, with in-

creasing time after exposure of cancers to prior therapy, there can be reduction in DNA hypermethylation and re-expression of transporters and other factors that have been downregulated as a consequence of exposure to chemotherapy and that may be required for drug efficacy.⁷⁶

Conversely, chemotherapy exposure can also in some cases result in recruitment of quiescent cells into active proliferation,⁷⁷ and there can be rapid regrowth of tumors between chemotherapy cycles, with increasingly rapid repopulation and resistance due to this “accelerated repopulation.”⁷⁸ Exposure of tumor cells to chemotherapy can also result in upregulation of expression of resistance factors.^{8,79,80} Tumors that regrow during chemotherapy or shortly after completion of chemotherapy are generally resistant to further chemotherapy,⁶ and this could potentially be attributable to the combination of accelerated repopulation⁷⁸ and upregulation of expression of resistance factors.^{8,79,80} In this instance, increasing drug doses might hypothetically somewhat improve therapy efficacy, but residual resistant quiescent cells would still ultimately limit outcome and flatten DRCs at highest doses.

Although the above could explain both DRC flattening and relapse patterns, it is stressed that our proposed explanation is highly speculative. In both preclinical and clinical studies, numerous resistance mechanisms have been noted in SCLC,^{8,60} and it remains unknown which ones are most important clinically. The field would benefit from carefully done translational studies (with tumor biopsies before therapy initiation and again at the time of maximum response and at the time of relapse) to assess how tumor cell growth characteristics, resistance factors, and factors required for therapy efficacy (e.g., drug obligate targets, drug transporters, drug activators, and proapoptotic factors) change over the course of therapy. If the initial presence of quiescent cells did prove to be central to the ultimate failure of therapy, then it would be particularly important to define potentially exploitable targets in these quiescent cells.




FIGURE 3. Weighted nonlinear meta-regression analysis (where the α parameter measures the relationship between dose and cell survival) or weighted linear regression analysis (where the β parameter measures the relationship between dose and cell survival) over higher doses (with elimination of data for doses of etoposide ≤ 100 , cisplatin ≤ 20 , epirubicin ≤ 20 , cyclophosphamide ≤ 200 , paclitaxel ≤ 35 , or ifosfamide ≤ 1250 mg/m²/wk or carboplatin \leq AUC 1.25/wk). Data for each of cisplatin ($n = 22$, $\alpha = -0.02$), carboplatin ($n = 6$, $\alpha = -0.39$), cyclophosphamide ($n = 3$, $\alpha = -0.01$), and ifosfamide ($n = 9$, $\alpha = -0.0004$) could be fitted to the nonlinear regression model, whereas data for each of etoposide ($n = 20$, $\beta = -0.09$), epirubicin ($n = 7$, $\beta = -0.48$), and paclitaxel ($n = 8$, $\beta = -0.11$) could not be fitted to the nonlinear regression model and were instead assessed by linear regression analysis.

The main purpose of this assessment was not to determine whether or not there is a benefit from administering higher doses of chemotherapy but rather to assess DRC shape, and the relatively small number of evaluable studies meant that statistical power was low. We did observe a downward slope in the DRCs for five of the seven agents over all dose intensities, and a downward slope was also seen for the two remaining drugs if the relatively flat early low-dose curve portion was omitted, but as noted above, the association between dose intensity and efficacy did not achieve statistical significance for any of the agents assessed. Recent reviews have also concluded that there is no proven benefit of higher dose intensity or dose density in SCLC.^{1,2,4,5,81} Nevertheless, available data suggest that it may be premature to conclude that dose is largely unimportant. In randomized trials, significant improvements in median survival,^{37,82–89} % 2-year survival,^{37,82,83,85–88} relapse-free survival,⁹⁰ complete response rates^{84,85,90–92} or overall response rates,^{52,83,87,89,93–95} or a statistically nonsignificant trend toward higher response

rates^{52,82,84,86,93,96} or longer survival^{90,97} have been reported with a variety of higher dose approaches or with addition of more agents to a therapy regimen. Only a minority of randomized trials have failed to show even a trend toward improvement in either survival or response with higher dose approaches or with increase in the number of agents used.^{45,48,98–103} Hence, despite the fact that even high drug doses are incapable of curing ED SCLC, it remains possible that higher drug doses may at least modestly improve palliation and short-term tumor control.

In summary, while maintaining relatively high doses of chemotherapy may possibly have a modestly positive impact in ED SCLC, DRC flattening at higher doses may preclude cure with currently available agents. The apparent DRC flattening would suggest that incurability of ED SCLC is due to deficiency or saturation of a factor or factors required for drug efficacy, and further progress will depend on our defining new therapeutic methods specifically targeting these resistant cells.

ACKNOWLEDGMENTS

Supported by NCI Cancer Center Support Grant 5-P30 CA16672 to UTMACC.

REFERENCES

- Crivellari G, Monfardini S, Stragliotto S, et al. Increasing chemotherapy in small-cell lung cancer: from dose intensity and density to megadoses. *Oncologist* 2007;12:79–89.
- Jackman DM, Johnson BE. Small-cell lung cancer. *Lancet* 2005;366:1385–1396.
- Mascaux C, Paesmans M, Berghmans T, et al. A systematic review of the role of etoposide and cisplatin in the chemotherapy of small cell lung cancer with methodology assessment and meta-analysis. *Lung Cancer* 2000;30:23–36.
- Kurup A, Hanna NH. Treatment of small cell lung cancer. *Crit Rev Oncol Hematol* 2004;52:117–126.
- Chua YJ, Steer C, Yip D. Recent advances in management of small-cell lung cancer. *Cancer Treat Rev* 2004;30:521–543.
- Huisman C, Postmus PE, Giaccone G, et al. Second-line chemotherapy and its evaluation in small cell lung cancer. *Cancer Treat Rev* 1999;25:199–206.
- Stewart DJ. Mechanisms of resistance to cisplatin and carboplatin. *Crit Rev Oncol Hematol* 2007;63:12–31.
- Stewart DJ. Tumor and host factors that may limit efficacy of chemotherapy in non-small cell and small cell lung cancer. *Crit Rev Oncol Hematol*. 2010;75:173–234.
- Stewart DJ, Raaphorst GP, Yau J, et al. Active vs. passive resistance, dose-response relationships, high dose chemotherapy, and resistance modulation: a hypothesis. *Invest New Drugs* 1996;14:115–130.
- Stewart DJ, Chiritescu G, Dahrouge S, et al. Chemotherapy dose-response relationships in non-small cell lung cancer and implied resistance mechanisms. *Cancer Treat Rev* 2007;33:101–137.
- Miller A, Hogestraeten B, Staquet M, et al. Reporting results of cancer treatment. *Cancer* 1981;47:207–214.
- Therasse P, Arbuck S, Eisenhauer E, et al. New guidelines to evaluate response to treatment in solid tumors. *J Natl Cancer Inst* 2000;92:205–216.
- Aisner J, Whitacre MY, Budman DR, et al. Cisplatin, doxorubicin, cyclophosphamide, and etoposide combination chemotherapy for small-cell lung cancer. *Cancer Chemother Pharmacol* 1992;29:435–438.
- Asamoto H, Kawahara M, Iwami F, et al. Cisplatin plus oral etoposide in the treatment of patients with advanced small cell lung cancer. Japan Clinical Oncology Group. *Jpn J Clin Oncol* 1998;28:745–748.
- Chen YM, Wu MF, Perng RP, et al. Phase II study of ifosfamide and etoposide chemotherapy for extensive-disease small-cell lung cancer. *Jpn J Clin Oncol* 1997;27:76–79.
- Clark PI, Slevin ML, Joel SP, et al. A randomized trial of two etoposide schedules in small-cell lung cancer: the influence of pharmacokinetics on efficacy and toxicity. *J Clin Oncol* 1994;12:1427–1435.
- Eckhardt S, Kolaric K, Vukas D, et al. Phase II study of 4'-epi-doxorubicin in patients with untreated, extensive small cell lung cancer. South-East European Oncology Group (SEEOG). *Med Oncol Tumor Pharmacother* 1990;7:19–23.
- Ettinger DS, Finkelstein DM, Sarma RP, et al. Phase II study of paclitaxel in patients with extensive-disease small-cell lung cancer: an Eastern Cooperative Oncology Group study. *J Clin Oncol* 1995;13:1430–1435.
- Fetscher S, Brugger W, Engelhardt R, et al. Standard- and high-dose etoposide, ifosfamide, carboplatin, and epirubicin in 107 patients with non-small-cell lung cancer: a mature follow-up report. *Ann Oncol* 1999;10:605–607.
- Frasci G, Nicoletta G, Comella P, et al. A weekly regimen of cisplatin, paclitaxel and topotecan with granulocyte-colony stimulating factor support for patients with extensive disease small cell lung cancer: a phase II study. *Br J Cancer* 2001;84:1166–1171.
- Gatzemeier U, Pawel JV, Laumen R, et al. Etoposide/vincristine-based chemotherapy with or without carboplatin in extensive-stage small cell lung cancer: a prospective randomized phase III trial. *Semin Oncol* 1994;21:31–35.
- Glisson BS, Kurie JM, Perez-Soler R, et al. Cisplatin, etoposide, and paclitaxel in the treatment of patients with extensive small-cell lung carcinoma. *J Clin Oncol* 1999;17:2309–2315.
- Glisson B, Lee JS, Palmer J, et al. Cisplatin, ifosfamide, and prolonged oral etoposide in the treatment of patients with extensive small cell lung carcinoma. *Cancer* 1998;82:301–308.
- Greco FA, Murphy PB, Hainsworth JD, et al. Prolonged administration of oral etoposide plus cisplatin in extensive stage small cell lung cancer. *Oncology* 1992;49(Suppl 1):34–38; discussion 39.
- Gridelli C, Perrone F, D'Aprile M, et al. Phase II study of intensive CEV (carboplatin, epirubicin and VP-16) plus G-CSF (granulocyte-colony stimulating factor) in extensive small cell lung cancer. *Eur J Cancer* 1995;31A:2424–2426.
- Gridelli C, Manzione L, Perrone F, et al. Carboplatin plus paclitaxel in extensive small cell lung cancer: a multicentre phase 2 study. *Br J Cancer* 2001;84:38–41.
- Gridelli C, Perrone F, Ianniello GP, et al. Carboplatin plus vinorelbine, a new well-tolerated and active regimen for the treatment of extensive-stage small-cell lung cancer: a phase II study. Gruppo Oncologico Centro-Sud-Isola. *J Clin Oncol* 1998;16:1414–1419.
- Gridelli C, Rossi A, Barletta E, et al. Carboplatin plus vinorelbine plus G-CSF in elderly patients with extensive-stage small-cell lung cancer: a poorly tolerated regimen. Results of a multicentre phase II study. *Lung Cancer* 2002;36:327–332.
- Grunberg SM, Crowley J, Hande KR, et al. Treatment of poor-prognosis extensive disease small-cell lung cancer with an all-oral regimen of etoposide and cyclophosphamide—a Southwest Oncology Group clinical and pharmacokinetic study. *Cancer Chemother Pharmacol* 1999;44:461–468.
- Hainsworth JD, Carrell D, Drengler RL, et al. Weekly combination chemotherapy with docetaxel and gemcitabine as first-line treatment for elderly patients and patients with poor performance status who have extensive-stage small cell lung carcinoma: a Minnie Pearl Cancer Research Network phase II trial. *Cancer* 2004;100:2437–2441.
- Hasturk S, Kurt B, Kocabas A, et al. Ifosfamide, cisplatin and etoposide (ICE) combined chemotherapy with recombinant human granulocyte colony-stimulating factor support in small cell lung cancer. *J Chemother* 1997;9:66–71.
- Higano CS, Crowley JJ, Veith RV, et al. A phase II trial of intravenous vinorelbine in previously untreated patients with extensive small cell lung cancer, a Southwest Oncology Group study. *Invest New Drugs* 1997;15:153–156.
- Johnson DH, Wolff SN, Hainsworth JD, et al. Extensive-stage small-cell bronchogenic carcinoma: intensive induction chemotherapy with high-dose cyclophosphamide plus high-dose etoposide. *J Clin Oncol* 1985;3:170–175.
- Kanitz E, Kolaric K, Jassem J, et al. Randomized phase II trial of high-dose 4'-epi-doxorubicin + cyclophosphamide versus high-dose 4'-epi-doxorubicin + cisplatin in previously untreated patients with extensive small cell lung cancer. *Oncology* 1992;49:327–332.
- Kirschling RJ, Grill JP, Marks RS, et al. Paclitaxel and G-CSF in previously untreated patients with extensive stage small-cell lung cancer: a phase II study of the North Central Cancer Treatment Group. *Am J Clin Oncol* 1999;22:517–522.
- Latreille J, Cormier Y, Martins H, et al. Phase II study of docetaxel (taxotere) in patients with previously untreated extensive small cell lung cancer. *Invest New Drugs* 1996;13:343–345.
- Loehrer PJ Sr, Ansari R, Gonin R, et al. Cisplatin plus etoposide with and without ifosfamide in extensive small-cell lung cancer: a Hoosier Oncology Group study. *J Clin Oncol* 1995;13:2594–2599.
- Luikart SD, Goutsou M, Mitchell ED, et al. Phase I/II trial of etoposide and carboplatin in extensive small-cell lung cancer. A report from the Cancer and Leukemia Group. *B. Am J Clin Oncol* 1993;16:127–131.
- Matsui K, Masuda N, Fukuoka M, et al. Phase II trial of carboplatin plus oral etoposide for elderly patients with small-cell lung cancer. *Br J Cancer* 1998;77:1961–1965.
- Mavroudis D, Pavlakou G, Blazoyiannakis G, et al. Sequential administration of cisplatin-etoposide followed by topotecan in patients with extensive stage small cell lung cancer. A multicenter phase II study. *Lung Cancer* 2003;39:71–76.
- Miller AA, Herndon JE II, Hollis DR, et al. Schedule dependency of 21-day oral versus 3-day intravenous etoposide in combination with

- intravenous cisplatin in extensive-stage small-cell lung cancer: a randomized phase III study of the Cancer and Leukemia Group B. *J Clin Oncol* 1995;13:1871–1879.
42. Mitchell EP, Perry MC, Luikart SD, et al. A phase I/II trial of etoposide and cisplatin in extensive small cell lung cancer: a cancer and leukemia group B study. *Lung Cancer* 1996;15:215–223.
 43. Murphy PB, Hainsworth JD, Greco FA, et al. A phase II trial of cisplatin and prolonged administration of oral etoposide in extensive-stage small cell lung cancer. *Cancer* 1992;69:370–375.
 44. Negoro S, Masuda N, Furuse K, et al. Dose-intensive chemotherapy in extensive-stage small-cell lung cancer. *Cancer Chemother Pharmacol* 1997;40(Suppl):S70–S73.
 45. Niell HB, Herndon JE II, Miller AA, et al. Randomized phase III intergroup trial of etoposide and cisplatin with or without paclitaxel and granulocyte colony-stimulating factor in patients with extensive-stage small-cell lung cancer: Cancer and Leukemia Group B Trial 9732. *J Clin Oncol* 2005;23:3752–3759.
 46. Noda K, Nishiwaki Y, Kawahara M, et al. Irinotecan plus cisplatin compared with etoposide plus cisplatin for extensive small-cell lung cancer. *N Engl J Med* 2002;346:85–91.
 47. Perez EA, Geoffroy FJ, Hillman S, et al. Phase II study of oral etoposide and intravenous paclitaxel in extensive-stage small cell lung cancer. *Lung Cancer* 2004;44:347–353.
 48. Pujol JL, Douillard JY, Riviere A, et al. Dose-intensity of a four-drug chemotherapy regimen with or without recombinant human granulocyte-macrophage colony-stimulating factor in extensive-stage small-cell lung cancer: a multicenter randomized phase III study. *J Clin Oncol* 1997;15:2082–2089.
 49. Saijo N, Horiike A. Topoisomerase I inhibitors in small-cell lung cancer. The Japanese experience. *Oncology (Williston Park)* 2004;18:11–16.
 50. Schiller JH, Kim K, Hutson P, et al. Phase II study of topotecan in patients with extensive-stage small-cell carcinoma of the lung: an Eastern Cooperative Oncology Group Trial. *J Clin Oncol* 1996;14:2345–2352.
 51. Schiller JH, Adak S, Cella D, et al. Topotecan versus observation after cisplatin plus etoposide in extensive-stage small-cell lung cancer: E7593—a phase III trial of the Eastern Cooperative Oncology Group. *J Clin Oncol* 2001;19:2114–2122.
 52. Sculier JP, Paesmans M, Lecomte J, et al. A three-arm phase III randomised trial assessing, in patients with extensive-disease small-cell lung cancer, accelerated chemotherapy with support of haematological growth factor or oral antibiotics. *Br J Cancer* 2001;85:1444–1451.
 53. Skarlos DV, Dimopoulos AM, Kosmidis P, et al. Docetaxel and gemcitabine combination, as first-line treatment, in patients with extensive disease small-cell lung cancer. A phase II study of the Hellenic Cooperative Oncology Group. *Lung Cancer* 2003;41:107–111.
 54. Socinski MA, Neubauer MA, Olivares J, et al. Phase II trial of paclitaxel, ifosfamide, and carboplatin in extensive-stage small cell lung cancer. *Lung Cancer* 2003;40:91–97.
 55. Sorensen BT, Nielsen K, Steenholdt S, et al. Treatment of extensive small cell lung cancer with carboplatin and teniposide. *Eur J Cancer* 1991;27:1109–1112.
 56. Takigawa N, Fujiwara K, Ueoka H, et al. Fractionated administration of irinotecan and cisplatin for treatment of extensive-disease small-cell lung cancer: a phase II study. *Anticancer Res* 2003;23:557–560.
 57. Trillet-Lenoir V, Soler P, Arpin D, et al. The limits of chemotherapy dose intensification using granulocyte colony stimulating factor alone in extensive small cell lung cancer. *Lung Cancer* 1996;14:331–341.
 58. Viren M, Liippo K, Ojala A, et al. Carboplatin and etoposide in extensive small cell lung cancer. *Acta Oncol* 1994;33:921–924.
 59. Stewart DJ, Nundy D, Maroun JA, et al. Bioavailability, pharmacokinetics, and clinical effects of an oral preparation of etoposide. *Cancer Treat Rep* 1985;69:269–273.
 60. Stewart DJ. Lung cancer resistance to chemotherapy. In DJ Stewart (Ed.), *Lung Cancer: Prevention, Management and Future Directions*. New York: Humana Press, 2010, pp 331–393.
 61. Banna GL, Simonelli M, Santoro A. High-dose chemotherapy followed by autologous hematopoietic stem-cell transplantation for the treatment of solid tumors in adults: a critical review. *Curr Stem Cell Res Ther* 2007;2:65–82.
 62. Dimanche-Boitrel MT, Garrido C, Chauffert B. Kinetic resistance to anticancer agents. *Cytotechnology* 1993;12:347–356.
 63. Mellor HR, Ferguson DJ, Callaghan R. A model of quiescent tumour microregions for evaluating multicellular resistance to chemotherapeutic drugs. *Br J Cancer* 2005;93:302–309.
 64. Jackson RC. The problem of the quiescent cancer cell. *Adv Enzyme Regul* 1989;29:27–46.
 65. Liang XJ, Shen DW, Gottesman MM. A pleiotropic defect reducing drug accumulation in cisplatin-resistant cells. *J Inorg Biochem* 2004;98:1599–1606.
 66. Shen DW, Su A, Liang XJ, et al. Reduced expression of small GTPases and hypermethylation of the folate binding protein gene in cisplatin-resistant cells. *Br J Cancer* 2004;91:270–276.
 67. Liang XJ, Mukherjee S, Shen DW, et al. Endocytic recycling compartments altered in cisplatin-resistant cancer cells. *Cancer Res* 2006;66:2346–2353.
 68. Roberson RS, Kussick SJ, Vallieres E, et al. Escape from therapy-induced accelerated cellular senescence in p53-null lung cancer cells and in human lung cancers. *Cancer Res* 2005;65:2795–2803.
 69. Ling YH, Zou Y, Perez-Soler R. Induction of senescence-like phenotype and loss of paclitaxel sensitivity after wild-type p53 gene transfection of p53-null human non-small cell lung cancer H358 cells. *Anticancer Res* 2000;20:693–702.
 70. Amaravadi RK, Yu D, Lum JJ, et al. Autophagy inhibition enhances therapy-induced apoptosis in a Myc-induced model of lymphoma. *J Clin Invest* 2007;117:326–336.
 71. Puig PE, Guilly MN, Bouchot A, et al. Tumor cells can escape DNA-damaging cisplatin through DNA endoreduplication and reversible polyploidy. *Cell Biol Int* 2008;32:1031–1043.
 72. Matsui K, Sawa T, Suzuki H, et al. Relapse of stage I small cell lung cancer ten or more years after the start of treatment. *Jpn J Clin Oncol* 2006;36:457–461.
 73. Sekine I, Nishiwaki Y, Kakinuma R, et al. Late recurrence of small-cell lung cancer: treatment and outcome. *Oncology* 1996;53:318–321.
 74. Johnson BE, Ihde DC, Bunn PA, et al. Patients with small-cell lung cancer treated with combination chemotherapy with or without irradiation. Data on potential cures, chronic toxicities, and late relapses after a five- to eleven-year follow-up. *Ann Intern Med* 1985;103:430–438.
 75. Vogelsang GB, Abeloff MD, Ettinger DS, et al. Long-term survivors of small cell carcinoma of the lung. *Am J Med* 1985;79:49–56.
 76. Stewart DJ, Issa JP, Kurzrock R, et al. Decitabine effect on tumor global DNA methylation and other parameters in a phase I trial in refractory solid tumors and lymphomas. *Clin Cancer Res* 2009;15:3881–3888.
 77. Durand RE, Vanderbyl SL. Schedule dependence for cisplatin and etoposide multifraction treatments of spheroids. *J Natl Cancer Inst* 1990;82:1841–1845.
 78. Davis AJ, Tannock JF. Repopulation of tumour cells between cycles of chemotherapy: a neglected factor. *Lancet Oncol* 2000;1:86–93.
 79. Liu SQ, Yu JP, Yu HG, et al. Activation of Akt and ERK signalling pathways induced by etoposide confer chemoresistance in gastric cancer cells. *Dig Liver Dis* 2006;38:310–318.
 80. Torigoe T, Izumi H, Ishiguchi H, et al. Cisplatin resistance and transcription factors. *Curr Med Chem Anticancer Agents* 2005;5:15–27.
 81. Berghmans T, Paesmans M, Lafitte JJ, et al. Role of granulocyte and granulocyte-macrophage colony-stimulating factors in the treatment of small-cell lung cancer: a systematic review of the literature with methodological assessment and meta-analysis. *Lung Cancer* 2002;37:115–123.
 82. Arriagada R, Le Chevalier T, Pignon JP, et al. Initial chemotherapeutic doses and survival in patients with limited small-cell lung cancer. *N Engl J Med* 1993;329:1848–1852.
 83. Buchholz E, Manegold C, Pilz L, et al. Standard versus dose-intensified chemotherapy with sequential reinfusion of hematopoietic progenitor cells in small cell lung cancer patients with favorable prognosis. *J Thorac Oncol* 2007;2:51–58.
 84. Masutani M, Ochi Y, Kadota A, et al. Dose-intensive weekly alternating chemotherapy for patients with small cell lung cancer: randomized trial, can it improve survival of patients with good prognostic factors? *Oncol Rep* 2000;7:305–310.
 85. Thatcher N, Girling DJ, Hopwood P, et al. Improving survival without

- reducing quality of life in small-cell lung cancer patients by increasing the dose-intensity of chemotherapy with granulocyte colony-stimulating factor support: results of a British Medical Research Council Multicenter Randomized Trial. Medical Research Council Lung Cancer Working Party. *J Clin Oncol* 2000;18:395–404.
86. Stewart WP, von Pawel J, Gatzemeier U, et al. Effects of granulocyte-macrophage colony-stimulating factor and dose intensification of V-ICE chemotherapy in small-cell lung cancer: a prospective randomized study of 300 patients. *J Clin Oncol* 1998;16:642–650.
 87. Fukuoka M, Masuda N, Negoro S, et al. CODE chemotherapy with and without granulocyte colony-stimulating factor in small-cell lung cancer. *Br J Cancer* 1997;75:306–309.
 88. Thatcher N, Qian W, Clark PI, et al. Ifosfamide, carboplatin, and etoposide with midcycle vincristine versus standard chemotherapy in patients with small-cell lung cancer and good performance status: clinical and quality-of-life results of the British Medical Research Council multicenter randomized LU21 trial. *J Clin Oncol* 2005;23:8371–8379.
 89. Pujol JL, Daures JP, Riviere A, et al. Etoposide plus cisplatin with or without the combination of 4'-epidoxorubicin plus cyclophosphamide in treatment of extensive small-cell lung cancer: a French Federation of Cancer Institutes multicenter phase III randomized study. *J Natl Cancer Inst* 2001;93:300–308.
 90. Humblet Y, Symann M, Bosly A, et al. Late intensification chemotherapy with autologous bone marrow transplantation in selected small-cell carcinoma of the lung: a randomized study. *J Clin Oncol* 1987;5:1864–1873.
 91. Johnson DH, Einhorn LH, Birch R, et al. A randomized comparison of high-dose versus conventional-dose cyclophosphamide, doxorubicin, and vincristine for extensive-stage small-cell lung cancer: a phase III trial of the Southeastern Cancer Study Group. *J Clin Oncol* 1987;5:1731–1738.
 92. O'Donnell MR, Ruckdeschel JC, Baxter D, et al. Intensive induction chemotherapy for small cell anaplastic carcinoma of the lung. *Cancer Treat Rep* 1985;69:571–575.
 93. Murray N, Livingston RB, Shepherd FA, et al. Randomized study of CODE versus alternating CAV/EP for extensive-stage small-cell lung cancer: an Intergroup Study of the National Cancer Institute of Canada Clinical Trials Group and the Southwest Oncology Group. *J Clin Oncol* 1999;17:2300–2308.
 94. Urban T, Chastang C, Lebas FX, et al. The addition of cisplatin to cyclophosphamide-doxorubicin-etoposide combination chemotherapy in the treatment of patients with small cell lung carcinoma: A randomized study of 457 patients. "Petites Cellules" Group. *Cancer* 1999;86:2238–2245.
 95. Sculier JP, Paesmans M, Bureau G, et al. Multiple-drug weekly chemotherapy versus standard combination regimen in small-cell lung cancer: a phase III randomized study conducted by the European Lung Cancer Working Party. *J Clin Oncol* 1993;11:1858–1865.
 96. Lorigan P, Woll PJ, O'Brien ME, et al. Randomized phase III trial of dose-dense chemotherapy supported by whole-blood hematopoietic progenitors in better-prognosis small-cell lung cancer. *J Natl Cancer Inst* 2005;97:666–674.
 97. Furuse K, Fukuoka M, Nishiwaki Y, et al. Phase III study of intensive weekly chemotherapy with recombinant human granulocyte colony-stimulating factor versus standard chemotherapy in extensive-disease small-cell lung cancer. The Japan Clinical Oncology Group. *J Clin Oncol* 1998;16:2126–2132.
 98. Ihde DC, Mulshine JL, Kramer BS, et al. Prospective randomized comparison of high-dose and standard-dose etoposide and cisplatin chemotherapy in patients with extensive-stage small-cell lung cancer. *J Clin Oncol* 1994;12:2022–2034.
 99. Ardzizoni A, Tjan-Heijnen VC, Postmus PE, et al. Standard versus intensified chemotherapy with granulocyte colony-stimulating factor support in small-cell lung cancer: a prospective European Organization for Research and Treatment of Cancer-Lung Cancer Group Phase III Trial-08923. *J Clin Oncol* 2002;20:3947–3955.
 100. Figueredo AT, Hryniuk WM, Strautmanis I, et al. Co-trimoxazole prophylaxis during high-dose chemotherapy of small-cell lung cancer. *J Clin Oncol* 1985;3:54–64.
 101. Mavroudis D, Papadakis E, Veslemes M, et al. A multicenter randomized clinical trial comparing paclitaxel-cisplatin-etoposide versus cisplatin-etoposide as first-line treatment in patients with small-cell lung cancer. *Ann Oncol* 2001;12:463–470.
 102. Miyamoto H, Kawakami Y, Arimoto T, et al. [Cisplatin etoposide versus cisplatin/etoposide/ifosfamide combination chemotherapy in small-cell lung cancer: a multicenter randomized controlled study. Hokkaido Study Group of Treatment for Small-Cell Lung Cancer]. *Gan To Kagaku Ryoho* 1991;18:1127–1134.
 103. Jett JR, Everson L, Therneau TM, et al. Treatment of limited-stage small-cell lung cancer with cyclophosphamide, doxorubicin, and vincristine with or without etoposide: a randomized trial of the North Central Cancer Treatment Group. *J Clin Oncol* 1990;8:33–38.